



PERFECT
SOLUTION
EDUCATION
GROUP

JC MATHEMATICS (H2) STUDY TIPS

**A USEFUL TECHNIQUE IN
VECTORS: THE SMART USE OF
EQUATING COEFFICIENTS**

PREPARED BY
Mr Jackie Lee



www.h2maths.com

JC_H2Maths_043



It is given that $\mathbf{a}$ and $\mathbf{b}$ are two non-zero and non-parallel vectors.

And if $\alpha\mathbf{a} + \beta\mathbf{b} = \lambda\mathbf{a} + \mu\mathbf{b} \Rightarrow \alpha\mathbf{a} - \lambda\mathbf{a} = \mu\mathbf{b} - \beta\mathbf{b} \Rightarrow (\alpha - \lambda)\mathbf{a} = (\mu - \beta)\mathbf{b}$,

hence, $\alpha - \lambda = 0$ and $\mu - \beta = 0 \Rightarrow \alpha = \lambda$ and $\beta = \mu$.

Example 11


Referred to the origin O , the points A and B are such that $\vec{OA} = \mathbf{a}$ and $\vec{OB} = \mathbf{b}$.

The point C lies on OA such that C is the mid-point of O and A . The point D lies on OB such that $OB:OD=3:2$. The lines AD and BC meet at the point E such that $BE = \lambda BC$ and $DE = \mu DA$.

(i) Find $\vec{OE}$ in terms of $\mathbf{a}$, $\mathbf{b}$ and λ .

(ii) Find $\vec{OE}$ in terms of $\mathbf{a}$, $\mathbf{b}$ and μ .

Hence, find the values of λ and μ .



$$(i) \vec{OE} = \frac{\lambda \vec{OC} + (1-\lambda) \vec{OB}}{\lambda + 1-\lambda}$$

$$= \lambda \left(\frac{1}{2} \mathbf{a} \right) + (1-\lambda) \mathbf{b}$$

$$(ii) \vec{OE} = \frac{\mu \vec{OA} + (1-\mu) \vec{OD}}{\mu + 1-\mu}$$

$$= \mu \mathbf{a} + (1-\mu) \left(\frac{2}{3} \mathbf{b} \right)$$

$$\therefore \frac{1}{2} \lambda = \mu \Rightarrow \lambda = 2\mu \quad (1)$$

$$1-\lambda = \frac{2}{3} (1-\mu) \quad (2)$$

$$\text{Sub. (1) into (2), } 3(1-2\mu) = 2(1-\mu)$$

$$3 - 6\mu = 2 - 2\mu$$

$$1 = 4\mu$$

$$\mu = \frac{1}{4} \quad (3)$$

$$\text{Sub. (3) into (1), } \lambda = 2\left(\frac{1}{4}\right) = \frac{1}{2}$$



PERFECT
SOLUTION
EDUCATION
GROUP

JC MATHEMATICS (H2)

STUDY TIPS

Explanation

The technique of equating coefficients relies on the property of non-parallel guide vectors. When a single position vector is expressed through two different geometric pathways using the same two base vectors, the two expressions must be identical.

Because the base vectors point in entirely different directions, they cannot substitute for or cancel one another. Consequently, the scalar multiplier for the first base vector on one side of the equation must equal its corresponding multiplier on the opposite side.

The same condition applies to the multipliers of the second base vector.

Matching these multipliers independently allows the vector components to be removed, converting a geometric relationship into a system of standard algebraic equations to resolve the unknowns.

This content is extracted from our lesson resources. Welcome to contact us for more tailor-made and in-depth learning.





SIGN UP NOW BEFORE IT'S TOO LATE



MATHS TUITION BY MR JACKIE LEE

Online • Face-to-Face • 1-to-1 • Group

- ✓ Former MOE Scholar
- ✓ Winner of International Mathematics Competitions
- ✓ PGDE Qualification (NIE) with **Distinction in Mathematics**
- ✓ Former **School Maths Teacher** in
 - Nan Hua High School
 - Raffles Institution
 - Anglo Chinese School (Indep)



Find out more about our
classes and courses at

www.h2maths.com



Scan to
WhatsApp



90705626